



Petal Fall

Fruit Set

Early Green Fruit

Fruit Colouring

Harvest

Post Harvest

Stage	Category	Treatment Options	Rate per Acre	Notes	
Petal Fall	Cherry Fruitworm	Rimon	540 – 800 mL	Make 1 st application @ petal fall. Repeat @ 10-14 day intervals. Max 3 apps per year. Most effective at egg hatch to 2 nd instar. PHI 8 days	
	Weevils	Actara 25WG	84-112 grams	Spray & residue are toxic to bees. PHI 3 Days	
	Botrytis	Or Captan Maestro	900 g	Mix together with Liberate spreader ¹ at 0.3-0.6 L/300 L water	
	Fertilizer	Terra Boost Boron Plus	1 L 0.5 L		
Fruit Set	Botrytis	Cantus	225 g	Mix together with Liberate spreader ¹ at 0.3 L / 300 L water	
	Fertilizer	Terra Boost	1 L		
	Aphids	Or Admire Assail	70 ml 23 – 34 g	If aphids are a problem, apply after flowering. Repeat 10 days later if needed. Apply in lots of water, 300-400 L / acre. PHI 3 days. Do not apply during bloom. PHI 7 Days. Also suppresses spanworm.	
Early Green Fruit	Fertilizer	TigerClaw Power-Cal Terra Boost Standard Premix	1 L 1 L 10 L	Tank mix and apply 10 days later.	
Fruit Colouring	SWD	Delegate	80 grams	PHI 3 Days	Do not use until registrations are confirmed for 2014. Berries become susceptible to SWD when fruit starts to colour.
		Entrust	40-53 grams	PHI 3 Days	
		Malathion 85E	400 mL/400L	PHI 2 Days	
		Pyganic	1 – 1.8 L	PHI 0 Days	
		Ripcord	60-70 mL	PHI 2 Days	
Anthracnose / Botrytis	Switch	390 g	Mix together with Liberate spreader ¹ at 0.3 L / 300 L water. (2 applications 7-10 days apart). PHI 1 Day		
Fertilizer	TigerClaw Fruit Finisher	2 L	Mix together with Liberate spreader ¹ at 0.3 L / 300 L water and apply. Repeat after each picking during harvest.		
Harvest	Fertilizer	TigerClaw Fruit Finisher	2 L	Mix together with Liberate spreader ¹ at 0.3 L / 300 L water and apply. Repeat after each picking during harvest.	

TerraLink Crop Programs are intended for use as an information resource only. Always confirm label rates. The label is the final authority

*Rates are based on mature plants using water volume of 300-400 L

¹ Liberate Spreader should be used at low rates when mixing with fertilizer.

Please check with your processor before applying pesticides because of varying maximum residue limits in countries that blueberries are shipped to.